

An affirmative answer to a problem posed by Zălinescu

Liangjin Yao*

November 9, 2009

Abstract

Recently, in [5] Zălinescu posed a question about the characterization of the intrinsic core of the Minkowski sum of two graphs associated with two maximal monotone operators. In this note we give an affirmative answer.

2000 Mathematics Subject Classification: Primary 47H05; Secondary 49J53, 52A41.

Keywords: Convex function, convex set, Fenchel conjugate, Fitzpatrick function, intrinsic core, maximal monotone operator, monotone operator, multifunction, relative algebraic interior, representative, set-valued operator.

1 Introduction

We suppose throughout this note that X is a real reflexive Banach space with norm $\|\cdot\|$ and dual product $\langle \cdot, \cdot \rangle$. We now introduce some notation. Let $A: X \rightrightarrows X^*$ be a *set-valued operator* or *multifunction* whose graph is defined by

$$\text{gra } A := \{(x, x^*) \in X \times X^* \mid x^* \in Ax\}.$$

The *domain* of A is $\text{dom } A := \{x \in X \mid Ax \neq \emptyset\}$. Recall that A is *monotone* if for all $(x, x^*), (y, y^*) \in \text{gra } A$ we have

$$\langle x - y, x^* - y^* \rangle \geq 0,$$

and A is *maximal monotone* if A is monotone and A has no proper monotone extension (in the sense of graph inclusions).

*Mathematics, Irving K. Barber School, UBC Okanagan, Kelowna, British Columbia V1V 1V7, Canada. E-mail: ljinyao@interchange.ubc.ca.

The *Fitzpatrick function* of A (see [1]) is given by

$$(1) \quad F_A : (x, x^*) \mapsto \sup_{(a, a^*) \in \text{gra } A} (\langle x, a^* \rangle + \langle a, x^* \rangle - \langle a, a^* \rangle).$$

For a function $f : X \rightarrow]-\infty, +\infty]$, the *domain* is $\text{dom } f := \{x \in X \mid f(x) < +\infty\}$ and $f^* : X^* \rightarrow [-\infty, +\infty] : x^* \mapsto \sup_{x \in X} (\langle x, x^* \rangle - f(x))$ is the *Fenchel conjugate* of f .

Given $F : X \times X^* \rightarrow]-\infty, +\infty]$, we say F is a *representative* of a maximal monotone operator A if F is lower semicontinuous and convex with $F \geq \langle \cdot, \cdot \rangle$, $F^* \geq \langle \cdot, \cdot \rangle$ and

$$\text{gra } A = \{(x, x^*) \mid F(x, x^*) = \langle x, x^* \rangle\}.$$

Following [2], it will be convenient to set $F^\Gamma : X^* \times X \rightarrow]-\infty, +\infty] : (x^*, x) \mapsto F(x, x^*)$, where $F : X \times X^* \rightarrow]-\infty, +\infty]$, and similarly for a function defined on $X^* \times X$.

We define $\widehat{F}$ (see [5]) by

$$\widehat{F}(x, x^*) := F(x, -x^*).$$

Let $a = (x, x^*), b = (y, y^*) \in X \times X^*$, we also set (see [4]) by

$$[a, b] = \langle x, y^* \rangle + \langle y, x^* \rangle.$$

Given a subset D of X , $\overline{D}$ is the *closure*, $\text{conv } D$ is the *convex hull*, and $\text{aff } D$ is the *affine hull*. The *conic hull* of D is denoted by $\text{cone } D := \{\lambda x \mid \lambda \geq 0, x \in D\}$. The *indicator function* $\iota_D : X \rightarrow]-\infty, +\infty]$ of D is defined by

$$x \mapsto \begin{cases} 0, & \text{if } x \in D; \\ +\infty, & \text{otherwise.} \end{cases}$$

The *intrinsic core* or *relative algebraic interior* of D , written as ${}^i D$ in [6], is

$${}^i D := \{a \in D \mid \forall x \in \text{aff}(D - D), \exists \delta > 0, \forall \lambda \in [0, \delta] : a + \lambda x \in D\}.$$

We define ${}^{ic} D$ by

$${}^{ic} D := \begin{cases} {}^i D, & \text{if } \text{aff } D \text{ is closed;} \\ \emptyset, & \text{otherwise.} \end{cases}$$

Zălinescu posed the following problem in [5]: Let $A, B : X \rightrightarrows X^*$ be maximal monotone. Is the implication

$${}^{ic} [\text{conv}(\text{gra } A - \text{gra}(-B))] \neq \emptyset \quad \Rightarrow \quad {}^{ic} [\text{dom } F_A - \text{dom } \widehat{F}_B] \neq \emptyset$$

true? Theorem 2.7 provides an affirmative answer to this question. It further shows that these two sets actually are equal.

2 Main result

Definition 2.1 (Fitzpatrick family) Let $A: X \rightrightarrows X^*$ be a maximal monotone operator. The associated Fitzpatrick family $\mathcal{F}_A$ consists of all functions $F: X \times X^* \rightarrow]-\infty, +\infty]$ that are lower semicontinuous and convex, and that satisfy $F \geq \langle \cdot, \cdot \rangle$, and $F = \langle \cdot, \cdot \rangle$ on $\text{gra } A$.

Fact 2.2 (Fitzpatrick) (See [1, Theorem 3.10].) Let $A: X \rightrightarrows X^*$ be a maximal monotone operator. Then for every $(x, x^*) \in X \times X^*$,

$$(2) \quad F_A(x, x^*) = \min \{F(x, x^*) \mid F \in \mathcal{F}_A\} \quad \text{and} \quad F_A^{*\top}(x, x^*) = \max \{F(x, x^*) \mid F \in \mathcal{F}_A\}.$$

Fact 2.3 (Simons) (See [4, Lemma 20.4(b)].) Let $A: X \rightrightarrows X^*$ be maximal monotone and $a := (x, x^*) \in X \times X^*$ with $\langle x, x^* \rangle = 0$. Suppose that there exists $u \in \mathbb{R}$ such that

$$[\text{gra } A, a] = \{u\}.$$

Then

$$[\text{dom } F_A, a] = \{u\}.$$

Theorem 2.4 Let $A, B: X \rightrightarrows X^*$ be maximal monotone. Then

$$(3) \quad \overline{\text{aff} [\text{gra } A - \text{gra}(-B)]} = \overline{\text{aff} [\text{dom } F_A - \text{dom } \widehat{F}_B]}.$$

Proof. We do and can suppose $(0, 0) \in \text{gra } A$ and $(0, 0) \in \text{gra } B$. We first show

$$(4) \quad [\text{dom } F_A - \text{dom } \widehat{F}_B] \subseteq \overline{\text{aff} [\text{gra } A - \text{gra}(-B)]}.$$

Suppose to the contrary that there exists $c \in X \times X^*$ such that $c \in [\text{dom } F_A - \text{dom } \widehat{F}_B]$ but $c \notin \overline{\text{aff} [\text{gra } A - \text{gra}(-B)]}$. By the Separation Theorem, there exist $a := (x, x^*) \in X \times X^*$ and $\delta \in \mathbb{R}$ such that

$$(5) \quad [a, c] > \delta > \sup \{[a, e] \mid e \in \overline{\text{aff} [\text{gra } A - \text{gra}(-B)]}\}.$$

Since $(0, 0) \in \text{gra } A$, $(0, 0) \in \text{gra } B$ and $\overline{\text{aff} [\text{gra } A - \text{gra}(-B)]}$ is a closed subspace, we have $\delta > 0$ and $[a, b - d] = 0, \forall b \in \text{gra } A, \forall d \in \text{gra}(-B)$. Thus,

$$(6) \quad [a, \text{gra } A] = \{0\} = [a, \text{gra}(-B)] = [(-x, x^*), \text{gra } B].$$

By $(0, 0) \in \text{gra } A$ and $(0, 0) \in \text{gra } B$ again,

$$(7) \quad F_A(a) = F_A(x, x^*) = 0, \quad F_B(-x, x^*) = 0.$$

Since $F_A(x, x^*) \geq \langle x, x^* \rangle$ and $F_B(-x, x^*) \geq \langle -x, x^* \rangle$, by (7), $\langle x, x^* \rangle = 0$. Thus by (6) and Fact 2.3,

$$[a, \text{dom } F_A] = \{0\} = [(-x, x^*), \text{dom } F_B] = [a, \text{dom } \widehat{F}_B].$$

Thus, $\lfloor a, \text{dom } F_A - \text{dom } \widehat{F}_B \rfloor = \{0\}$, which contradicts (5). Hence

$$(8) \quad \left[\text{dom } F_A - \text{dom } \widehat{F}_B \right] \subseteq \overline{\text{aff} [\text{gra } A - \text{gra}(-B)]}.$$

And thus $\overline{\text{aff} [\text{dom } F_A - \text{dom } \widehat{F}_B]} \subseteq \overline{\text{aff} [\text{gra } A - \text{gra}(-B)]}$. Hence

$$\overline{\text{aff} [\text{gra } A - \text{gra}(-B)]} = \overline{\text{aff} [\text{dom } F_A - \text{dom } \widehat{F}_B]}.$$

■

Fact 2.5 (Zălinescu) (See [5, Lemma 2 and Theorem 3].) *Let $A, B : X \rightrightarrows X^*$ be maximal monotone, and let F_1, F_2 be representatives of A, B , respectively. Then*

$${}^{ic} [\text{gra } A - \text{gra}(-B)] = {}^{ic} [\text{conv}(\text{gra } A - \text{gra}(-B))]$$

and

$${}^{ic} [\text{dom } F_1 - \text{dom } \widehat{F}_2] \subseteq [\text{gra } A - \text{gra}(-B)] \subseteq \text{conv} [\text{gra } A - \text{gra}(-B)] \subseteq [\text{dom } F_1 - \text{dom } \widehat{F}_2].$$

If ${}^{ic} [\text{dom } F_1 - \text{dom } \widehat{F}_2] \neq \emptyset$, then

$$(9) \quad {}^{ic} [\text{dom } F_1 - \text{dom } \widehat{F}_2] = {}^{ic} [\text{gra } A - \text{gra}(-B)] = {}^{ic} [\text{conv}(\text{gra } A - \text{gra}(-B))].$$

Remark 2.6 If X is finite-dimensional, the intrinsic core of a convex set $D \subseteq X$ is the same as the relative interior of D in the sense of Rockafellar[3]. Then ${}^{ic} [\text{dom } F_1 - \text{dom } \widehat{F}_2] = {}^i [\text{dom } F_1 - \text{dom } \widehat{F}_2] \neq \emptyset$ by [3, Theorem 6.2]. Thus, (9) always holds.

Our main result comes the following which provides an affirmative answer to the question posed by Zălinescu.

Theorem 2.7 *Let $A, B : X \rightrightarrows X^*$ be maximal monotone such that ${}^{ic} [\text{conv} (\text{gra } A - \text{gra}(-B))] \neq \emptyset$. Then*

$${}^{ic} [\text{gra } A - \text{gra}(-B)] = {}^{ic} [\text{conv} (\text{gra } A - \text{gra}(-B))] = {}^{ic} [\text{dom } F_A - \text{dom } \widehat{F}_B].$$

Moreover, if F_1, F_2 are representatives of A, B , respectively, then

$$(10) \quad {}^{ic} [\text{dom } F_1 - \text{dom } \widehat{F}_2] = {}^{ic} [\text{conv} (\text{gra } A - \text{gra}(-B))] = {}^{ic} [\text{dom } F_A - \text{dom } \widehat{F}_B].$$

Proof. Let $a \in {}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))]$. Then we have $a \in [\text{dom } F_A - \text{dom } \widehat{F}_B]$ and $\text{cone}[\text{conv}(\text{gra } A - \text{gra}(-B)) - a]$ is a closed subspace. By Theorem 2.4,

$$\begin{aligned} \text{cone}[\text{conv}(\text{gra } A - \text{gra}(-B)) - a] &\subseteq \text{cone}[\text{dom } F_A - \text{dom } \widehat{F}_B - a] \\ &\subseteq \text{aff}[\text{dom } F_A - \text{dom } \widehat{F}_B - a] = \text{aff}[\text{dom } F_A - \text{dom } \widehat{F}_B] - \{a\} \\ &\subseteq \overline{\text{aff}[\text{gra } A - \text{gra}(-B)] - \{a\}} \subseteq \overline{\text{aff}[\text{gra } A - \text{gra}(-B) - a]} \\ &= \overline{\text{aff}[\text{conv}(\text{gra } A - \text{gra}(-B)) - a]} \subseteq \text{cone}[\text{conv}(\text{gra } A - \text{gra}(-B)) - a]. \end{aligned}$$

Hence $\text{cone}[\text{dom } F_A - \text{dom } \widehat{F}_B - a] = \text{cone}[\text{conv}(\text{gra } A - \text{gra}(-B)) - a]$ is a closed subspace. Thus $a \in {}^{ic}[\text{dom } F_A - \text{dom } \widehat{F}_B]$. By Fact 2.5,

$${}^{ic}[\text{gra } A - \text{gra}(-B)] = {}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))] = {}^{ic}[\text{dom } F_A - \text{dom } \widehat{F}_B].$$

And by Fact 2.2,

$$\text{conv}[\text{gra } A - \text{gra}(-B)] \subseteq [\text{dom } F_1 - \text{dom } \widehat{F}_2] \subseteq [\text{dom } F_A - \text{dom } \widehat{F}_B].$$

Similar to the proof above, see that (10) holds. ■

Theorem 2.8 *Let $A, B : X \rightrightarrows X^*$ be maximal monotone, and F_1, F_2 be representatives of A, B , respectively. Then*

$$(11) \quad {}^{ic}[\text{dom } F_1 - \text{dom } \widehat{F}_2] = {}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))] = {}^{ic}[\text{gra } A - \text{gra}(-B)].$$

Proof. We consider two cases.

Case 1: ${}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))] = \emptyset$. Assume that ${}^{ic}[\text{dom } F_1 - \text{dom } \widehat{F}_2] \neq \emptyset$. Then by Fact 2.5, ${}^{ic}[\text{dom } F_1 - \text{dom } \widehat{F}_2] = {}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))] = \emptyset$. This is a contradiction.

Case 2: ${}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))] \neq \emptyset$. Apply Theorem 2.7.

Combining the above results, we see that (11) holds. ■

Corollary 2.9 *Let $A, B : X \rightrightarrows X^*$ be maximal monotone, and F_1, F_2 be representatives of A, B , respectively. Assume ${}^{ic}[\text{conv}(\text{gra } A - \text{gra}(-B))] \neq \emptyset$. Then*

$$(12) \quad \overline{[\text{dom } F_1 - \text{dom } \widehat{F}_2]} = \overline{\text{conv}[(\text{gra } A - \text{gra}(-B))]} = \overline{[\text{gra } A - \text{gra}(-B)]}.$$

In particular,

$$\overline{[\text{dom } F_A - \text{dom } \widehat{F}_B]} = \overline{\text{conv}[(\text{gra } A - \text{gra}(-B))]} = \overline{[\text{gra } A - \text{gra}(-B)]}.$$

Proof. Given a convex set $D \subseteq X$, assume that ${}^{ic}D \neq \emptyset$, then $\overline{{}^{ic}D} = \overline{D}$. By Theorem 2.8,

$$\overline{[\text{dom } F_1 - \text{dom } \widehat{F}_2]} = \overline{\text{conv}[(\text{gra } A - \text{gra}(-B))]} = \overline{{}^{ic}[\text{gra } A - \text{gra}(-B)]} \subseteq \overline{[\text{gra } A - \text{gra}(-B)]}$$

Hence (12) holds. ■

Acknowledgment

The author thanks Heinz Bauschke and Xianfu Wang for valuable discussions.

References

- [1] S. Fitzpatrick, “Representing monotone operators by convex functions”, in *Workshop/Miniconference on Functional Analysis and Optimization (Canberra 1988)*, Proceedings of the Centre for Mathematical Analysis, Australian National University, vol. 20, Canberra, Australia, pp. 59–65, 1988.
- [2] J.-P. Penot, “The relevance of convex analysis for the study of monotonicity”, *Nonlinear Analysis*, vol. 58, pp. 855–871, 2004.
- [3] R.T. Rockafellar, *Convex Analysis*, Princeton University Press, 1970.
- [4] S. Simons, *From Hahn-Banach to Monotonicity*, Springer-Verlag, 2008.
- [5] C. Zălinescu, “A new convexity property for monotone operators”, *Journal of Convex Analysis*, vol. 13, pp. 883–887, 2006.
- [6] C. Zălinescu, *Convex Analysis in General Vector Spaces*, World Scientific Publishing, 2002.